

POSTER PRESENTATION

Open Access

Indoleamine 2,3-dioxygenase is a critical resistance mechanism in anti-tumor T cell immunotherapy targeting CTLA-4

Rikke B Holmgaard^{1*}, Dmitriy Zamarin¹, David H Munn³, Jedd D Wolchok¹, James P Allison^{1,2}

From Society for Immunotherapy of Cancer 28th Annual Meeting
National Harbor, MD, USA. 8-10 November 2013

The CTLA-4 blocking antibody ipilimumab results in durable responses in metastatic melanoma, though therapeutic benefit has been limited to a fraction of patients. This calls for identification of resistance mechanisms and development of combinatorial strategies. We examine the inhibitory role of indoleamine 2,3-dioxygenase (IDO) on the anti-tumor efficacy of CTLA-4 blockade. In IDO knockout mice treated with anti-CTLA-4 antibody, we demonstrate a striking delay in B16 melanoma tumor growth and increased overall survival when compared to wild type mice. To highlight the therapeutic relevance of these findings, we show that CTLA-4 blockade strongly synergizes with IDO inhibitors to mediate rejection of both IDO-expressing and non-expressing poorly immunogenic tumors, emphasizing the importance of the inhibitory role of both tumor- and host-derived IDO. This effect was T-cell dependent, leading to enhanced infiltration of tumor-specific effector T cells and a marked increase in the effector to regulatory T cell ratios in the tumors. Overall, these data demonstrate the immunosuppressive role of IDO in the context of anti-CTLA-4 immunotherapy and provide a strong incentive to clinically explore combination therapies utilizing IDO inhibitors irrespective of IDO expression by the tumor cells.

Authors' details

¹Department of Immunology, Memorial Sloan-Kettering Cancer Center, New York, NY, USA. ²Department of Immunology, MD Anderson Cancer Center, Houston, TX, USA. ³Cancer Center and Department of Pediatrics, Georgia Regents University, Augusta, GA, USA.

Published: 7 November 2013

¹Department of Immunology, Memorial Sloan-Kettering Cancer Center, New York, NY, USA

Full list of author information is available at the end of the article

doi:10.1186/2051-1426-1-S1-P77

Cite this article as: Holmgaard et al.: Indoleamine 2,3-dioxygenase is a critical resistance mechanism in anti-tumor T cell immunotherapy targeting CTLA-4. *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):P77.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

